

GROUP	1	B	INSECTICIDE
-------	---	---	-------------

FIFRA 24(c) REGISTRATION
EPA Reg. No. 10163-169
EPA SLN No. WA-010019
Expires: December 31, 2014

**REGISTRATION FOR SPECIAL LOCAL NEED
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON
PRE-BLOOM APPLICATION OF IMIDAN 70-W
IN GRAPES TO CONTROL GRAPE MEALYBUG**

This label for Imidan expires and must not be distributed or used in accordance with this SLN registration after December 31, 2014.

Imidan® 70-W

Agricultural Insecticide

EPA Reg. No. 10163-169

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label.
- This labeling must be in the possession of the user at the time of pesticide application.
- For use under the SLN label do not apply this product through any type of irrigation system.

CROP

Grapes

PESTS CONTROLLED

Grape Mealybug, Cutworms

USE RATE

Pre-bloom application only – 3 lbs Imidan 70-W (2 lb ai) per acre.

METHOD OF APPLICATION

For pre-bloom applications only, apply up to 3 lbs Imidan 70-W (2 lb ai) in combination with oil or spreader sticker. Use sufficient volume for adequate coverage.

RESTRICTIONS / PRECAUTIONS

- Do not apply more than 3 times per season
- For **groundboom and airblast applications** during the **growing season**, do not apply this product within 25 feet of (1) buildings occupied by humans for residential, commercial, or business purposes, including but not limited homes, farmworker housing, or other residential buildings, schools, daycare centers, nursing homes, hospitals, and (2) outdoor recreational areas such as school grounds, athletic fields, and parks. Non-residential agricultural buildings, including barns, livestock facilities, sheds, and outhouses are not included in this prohibition.
- Aerial application is prohibited.
- This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or broadleaf weeds. Do not apply this product or allow it to drift to blooming broadleaf weeds. Broadleaf weed bloom must be controlled prior to the application in vineyards with 5 or more blooms per square yard
- This pesticide is extremely toxic to fish. IMIDAN 70-W should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.
- Apply as a pre-bloom spray only

WSDA CONTAINER DISPOSAL GUIDANCE

- Completely empty the contents of the paper container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

NOTE

This product is sold subject to the CONDITIONS OF SALE set forth on the container label.

24(c) REGISTRANT

Gowan Company
P.O. Box 5569
Yuma, AZ 85366

IMIDAN[®] is a registered trademark of Gowan Company, L.L.C
Rev. 3-23-10